

Direct equivalence between quantum phase transition phenomena in radiation-matter and magnetic systems: scaling of entanglement

J. RESLEN¹(*), L. QUIROGA^(**)¹ and N. F. JOHNSON²(***)

¹ *Departamento de Física, Universidad de Los Andes, A.A.4976, Bogotá D.C., Colombia*

² *Centre for Quantum Computation and Department of Physics, University of Oxford, Clarendon Laboratory, Parks Road, OX1 3PU, U.K*

PACS. 03.65.Ud – Entanglement and quantum nonlocality.

PACS. 73.43.Nq – Quantum phase transitions.

PACS. 75.10.-b – General theory and models of magnetic ordering.

Abstract. – We show that the quantum phase transition arising in a standard radiation-matter model (Dicke model) belongs to the same universality class as the infinitely-coordinated, transverse field XY model. The effective qubit-qubit exchange interaction is shown to be proportional to the square of the qubit-radiation coupling. A universal finite-size scaling is derived for the corresponding two-qubit entanglement (concurrence) and a size-consistent effective Hamiltonian is proposed for the qubit subsystem.

Quantum phase transitions (QPTs) are associated with a dramatic change in the physical properties of a system at zero temperature when a parameter varies around its critical value. It is well-known that very different systems can exhibit similar behavior in this critical regime, giving rise to the concept of *universality*. Enlarging a given universality class by the addition of systems from very different areas of physics, is a very important step toward unifying our understanding of the basic physics underlying apparently disconnected complex phenomena. Recently there have been studies of light-controlled condensed matter systems displaying QPTs with atoms in extreme one-dimensional confinements [1], ions driven by properly tuned and pulsed light [2] and fermionic atoms in optical superlattices [3]. Fully quantum mechanical models of radiation-matter systems are also being considered, and are important for several reasons: Scalable and distributed quantum information processing (QIP) devices will demand the integration of matter quantum bits (qubits) such as atoms, trapped ions, semiconductor quantum dots or SQUIDS with photons. In addition, the capability of photons to control and modify the coupling between physically distant qubits makes them appropriate for manipulating and transferring quantum information.

(*) E-mail: j-reslen@uniandes.edu.co

(**) E-mail: lquirola@uniandes.edu.co

(***) E-mail: n.johnson@physics.ox.ac.uk

QPTs themselves are beginning to be studied using concepts from quantum information theory. Special attention has been paid to the connections between subsystem entanglement and critical phenomena [4–6]. There is an obvious correspondence between condensed matter magnetic systems comprising N spin-1/2 particles, and quantum registers with N qubits. Most of the quantum information-based analysis on QPTs performed to date corresponds to critical magnetic systems where the exchange coupling between spins is fixed from the outset by Nature. However there is also interest in studying QPTs where the spin/qubit couplings can be controlled externally, using light for example. One of the most famous radiation-matter systems giving rise to a QPT is the Dicke model [7, 8]. In this model, N two-level systems (qubits) are coupled to a single radiation mode (cavity). Although no direct coupling exists between qubits, a superradiant phase arises as the qubit-cavity coupling strength increases. Quantum chaos and entanglement in the Dicke model have been studied recently using spin-boson transformation techniques [9, 10]. However a full understanding of the entanglement (concurrence) finite-size scaling behavior and the associated universality class, are still lacking.

Here we show that the reduced dynamics of a qubit system embedded in a photon field, as described by the Dicke model, displays magnetic-like behavior and corresponds to the same universality class of quantum critical phenomena as an infinitely-coordinated ferromagnetic system. The qubit-cavity detuning determines an effective magnetic field while the square of the qubit-cavity coupling strength yields the effective qubit-qubit ‘exchange’ interaction. As a result, we are able to establish an unambiguous quantitative connection between the associated scaling exponents and effective qubit Hamiltonian in two quite different physical systems. Although Lambert et al. [10] have already noted similarities between the Dicke and collective XY model in terms of scaling exponents for the concurrence, the present work establishes a systematic link between these models and lays these similarities on a firm theoretical footing.

We are interested in a system formed by N identical two-level atoms (qubits) coupled to a single-mode cavity, described by the Hamiltonian

$$H = a^\dagger a + \frac{\epsilon}{2} \sum_{i=1}^N \sigma_{i,z} + \frac{\lambda}{\sqrt{N}} \sum_{i=1}^N (a^\dagger + a)(\sigma_i^\dagger + \sigma_i) \quad (1)$$

where $a^\dagger$ (a) is the creation (annihilation) operator for single-mode cavity photons, $\sigma_{i,z} = |1_i\rangle\langle 1_i| - |0_i\rangle\langle 0_i|$, $\sigma_i^\dagger = |1_i\rangle\langle 0_i|$, $\sigma_i = |0_i\rangle\langle 1_i|$ with $|0_i\rangle$ and $|1_i\rangle$ ($i = 1, 2, \dots, N$) being the ground and excited states of the i ’th qubit, respectively, and λ is the qubit-cavity coupling. Note that we do not make a rotating wave approximation. We take the energy of a confined photon as the unit of energy. Resonance between qubit energy levels and photons corresponds to $\epsilon = 1$. This Hamiltonian can also be written as

$$H = a^\dagger a + \epsilon J_z + \frac{2\lambda}{\sqrt{N}} (a^\dagger + a) J_x \quad (2)$$

where $J_z = \frac{1}{2} \sum_{i=1}^N \sigma_{i,z}$ and $J_x = \frac{1}{2} \sum_{i=1}^N (\sigma_i^\dagger + \sigma_i)$. We denote this latter Hamiltonian as $H = H_0 + V$ where $H_0 = a^\dagger a$ and $V = \epsilon J_z + \frac{2\lambda}{\sqrt{N}} (a^\dagger + a) J_x$.

We shall be concerned with physical qubit quantities such as ‘magnetization’ and entanglement. These quantities can be represented by a generic qubit operator Q . Hence in thermodynamic equilibrium at inverse temperature $\beta = (k_B T)^{-1}$ any qubit expectation value can be expressed as

$$\langle Q \rangle = \frac{\text{Tr}_{qb}[Q \text{Tr}_{ph}[e^{-\beta H}]]}{\text{Tr}_{qb}[\text{Tr}_{ph}[e^{-\beta H}]]} = \frac{\text{Tr}_{qb}[Q e^{-\beta H_{qb}}]}{\text{Tr}_{qb}[e^{-\beta H_{qb}}]} \quad (3)$$

where Tr_{qb} (Tr_{ph}) denotes a trace over the qubit (cavity) subsystem and the last equality is the formal definition of an effective qubit Hamiltonian H_{qb} . In order to eliminate the cavity degrees of freedom and obtain a size-consistent H_{qb} (i.e. a reduced qubit Hamiltonian which scales with the system's size) we follow the procedure sketched out by Polatsek and Becker [11]. The resulting effective qubit Hamiltonian can be written as

$$H_{qb} = -\frac{1}{\beta} \ln \langle e^{-\beta(L_0+V)} \rangle_{ph} \quad (4)$$

where the thermal averaging is carried out with respect to the cavity degrees of freedom, i.e. $\langle O \rangle_{ph} = Tr_{ph}(Oe^{-\beta a^\dagger a})/Tr_{ph}(e^{-\beta a^\dagger a})$; L_0 is the Liouville operator associated with H_0 and defined by $L_0 O = [a^\dagger a, O]$ for any operator O . A cumulant expansion is well-suited to calculating a size-consistent H_{qb} , yielding

$$H_{qb} = -\frac{1}{\beta} \sum_{n=1}^{\infty} (-1)^n \frac{\beta^n}{n!} H_n \quad (5)$$

where H_n represents the qubit-operator corresponding to the n 'th cumulant. Cumulants coming from eq. (4) are themselves operators which act on qubit states, hence any expression involving cumulants must be properly symmetrized. The lowest-order cumulants are:

$$H_1 = \epsilon J_z \quad , \quad H_2 = \left[\frac{2\lambda}{\sqrt{N}} \right]^2 (2h(\beta) + 1) J_x^2 \quad (6)$$

where $h(\beta) = (e^\beta - 1)^{-1}$ is the Bose factor which determines the average photon number in an isolated cavity at temperature T . Due to the superradiance phenomenon which occurs in the Dicke model, the actual photon number has very different behavior on either side of the quantum and/or thermal phase transition. It can be seen that H_1 corresponds to the non-interacting qubit system while H_2 gives rise to the lowest order effective qubit-qubit coupling.

Every cumulant contains a term of the form

$$\langle V L_0^n V \rangle_{ph} = \left[\frac{2\lambda}{\sqrt{N}} \right]^2 [(1 + (-1)^n)h(\beta) + 1] J_x^2 \quad (7)$$

which enables us to perform a closed-form summation in eq. (5) over an infinite set of contributing terms. Up to second order in V , this renormalization summation yields a temperature-dependent effective Hamiltonian given by

$$H_{qb}^{(2)}(\beta) = \epsilon J_z - \left[\frac{2\lambda}{\sqrt{N}} \right]^2 \left[1 + \frac{2}{\beta(h(\beta) + 1)} \right] J_x^2 \quad (8)$$

At zero temperature (i.e. $\beta \rightarrow \infty$) this effective Hamiltonian is a special case of that studied in ref. [6] corresponding to a ferromagnetic infinitely-coupled XY model in a transverse magnetic field with a QPT at $2\lambda_c = 1$ in the thermodynamic limit (i.e. $N \rightarrow \infty$). Our analysis therefore provides a novel demonstration of the well-known QPT arising in the Dicke model at $\lambda_c = 1/2$ [8]. The broken symmetry associated with the phase transition is related to parity for the Dicke model as given by $e^{i\pi(a^\dagger a + J_z + \frac{N}{2})}$, while for the effective $H_{qb}^{(2)}$ model this becomes $e^{i\pi(J_z + \frac{N}{2})}$. In contrast to a previously employed continuous variable representation [9, 10], this effective anisotropic XY model Hamiltonian $H_{qb}^{(2)}$ provides an ideal starting point

TABLE I – Critical exponent b at zero temperature for the finite- N scaling of the x -magnetization at λ_c (first row), z -magnetization at λ_c (second row), difference between the concurrences at λ_c for the thermodynamic limit and for N qubits (third row), difference between the maxima of the concurrences for the thermodynamic limit and for N qubits (fourth row), and the difference between the coupling strength for N qubits at maximum concurrence and λ_c (fifth row). $N \leq 35$ for the Dicke model and $100 \leq N \leq 500$ for the anisotropic XY model.

$f(N) \sim N^b$	Dicke	XY
$\frac{\sqrt{\langle J_z^2 \rangle_{\lambda_c}}}{N}$	-0.35 ± 0.01	-0.34 ± 0.01
$\frac{1}{2} - \frac{\langle J_z \rangle_{\lambda_c}}{N}$	-0.54 ± 0.01	-0.55 ± 0.01
$C_\infty(\lambda_c) - C_N(\lambda_c)$	-0.26 ± 0.01	-0.30 ± 0.01
$C_\infty^M - C_N^M$	-0.28 ± 0.03	-0.30 ± 0.03
$\lambda_N^M - \lambda_c$	-0.65 ± 0.03	-0.66 ± 0.03

for studying the QPT of the Dicke model since we can follow the qubit subsystem's quantum-mechanical evolution as a function of the effective qubit-qubit coupling. Furthermore, this mapping onto the infinitely-coordinated XY model which results from our approach, yields a deep understanding of the scaling properties for finite N as well as the mean-field behavior of the Dicke model in the thermodynamic limit. Fig. 1 shows the ground-state energy (at zero




Fig. 1




Fig. 2

Fig. 1 – Ground state energy and z -magnetization (inset) as a function of λ for $N = 15$ qubits, as obtained from the full Dicke Hamiltonian ($\times$) and from the effective Hamiltonian ($\circ$).

Fig. 2 – Non-rescaled concurrence as a function of λ for the XY model and for the full Dicke Hamiltonian (inset) for $N=15$ and 35 .

temperature) of a Dicke system with N qubits, as a function of the qubit-radiation coupling λ . For simplicity, we only consider the case of resonance from now on, i.e. $\epsilon = 1$. For comparison, the ground-state energy produced by the effective Hamiltonian in eq. (8) is also included in the plot. The agreement between both ground-state energies is excellent even for low N . The qubit subsystem's ‘ z -magnetization’, which is the order parameter for this QPT, undergoes a quantum phase transition at $\lambda_c = 1/2$ as seen in fig. 1 (see inset) for both the exact Dicke model and the effective Hamiltonian. This indicates that the physics of the QPT in the Dicke




Fig. 3




Fig. 4

Fig. 3 – Dependence of $\lambda_N^M - \lambda_c$ (lower curves) and $C_\infty^M - C_N^M$ (upper curves) on N for H ($\triangle$) and $H_{qb}^{(2)}$ ($\times$). λ_N^M is the value of λ for which the concurrence is maximum, while C_N^M is the maximum of the rescaled concurrence.

Fig. 4 – Finite-size scaling for the rescaled concurrence in the Dicke model. Points corresponding to different N collapse on the same curve. Inset: Finite-size scaling for the x-magnetization $m_x = \frac{\sqrt{\langle J_x^2 \rangle}}{N}$.

model is indeed captured by the effective Hamiltonian of eq. (8).

As a more stringent test of the effective Hamiltonian in eq. (8), we now compare other thermodynamic as well as quantum features. An important ground-state quantum feature is the entanglement between two qubits. We adopt the concurrence C_{NR} as a measure of this entanglement [12]. We also study the *rescaled concurrence* $C = NC_{NR}$ in order to obtain non-trivial information in the thermodynamic limit where $C_{NR} = 0$. In contrast with thermodynamic results that only involve the calculation of the partition function and averages obtained from it, the concurrence probes the internal structure of the ground-state wavefunction in a more detailed way. In fig. 2 we display the evolution of the concurrence as a function of λ , for both the exact Dicke model and the effective Hamiltonian of eq. (8). The qualitative form of the concurrence for both models is very similar, even though the effective Hamiltonian overestimates the maximum value. This overestimation is easily understood: the effective Hamiltonian drops higher-order interaction terms which incorporate additional interactions among the qubits and would hence modify the ground-state wavefunction. In the thermodynamic limit, the rescaled concurrence displays a singularity at $\lambda = \lambda_c$ in agreement with the findings in refs. [6, 10]. However, at λ_c the rescaled concurrence goes to 1 for the XY model while it goes to $1 - \frac{\sqrt{2}}{2}$ for the Dicke model.

As a final test of the reduced Hamiltonian of eq. (8), we consider the finite-size scaling of the ‘magnetization’ as well as the concurrence. Results can be compared with those reported in ref. [13] and show excellent agreement between the predictions of the Dicke model and the effective Hamiltonian.

At zero temperature, the energy and the z-magnetization are identical for both systems in the thermodynamic limit. Physical quantities depending on the ground state structure, such as the x-magnetization and concurrence, display some differences. However, the scaling exponents agree for both models as can be seen in fig. 3 and table I. Even for small N in the Dicke model, the quantities $\lambda_N^M - \lambda_c$ and $C_\infty^M - C_N^M$ can be seen to lie on well-defined straight lines. For the XY model, large N results lie on straight lines which are parallel to the Dicke

model results. The Dicke model results approach the thermodynamic limit much faster than those for the XY model. This is directly related to the fact that terms have been thrown away in eq. (8) which include interactions between three and more spins, thereby enhancing the mean-field nature of this system.

The excellent agreement between the critical exponents for the two models provides clear evidence that they both belong to the same universality class. It is well known that in both Hamiltonians, each qubit is coupled to every other qubit with the same strength and hence no typical lengthscale can be associated with them. Botet et al. [13] proposed relating the infinitely-coordinated magnetic system to short-range interaction models, and hence were able to perform a systematic study of finite-size scaling functions for the x-magnetization and gap energy in the case of a Hamiltonian such as $H_{qb}^{(2)}$. Following this line of reasoning, we show in fig. 4 that this scaling hypothesis is valid for the Dicke model as well. In particular there exist scaling functions in the critical region $|\lambda - \lambda_c| \rightarrow 0$: the concurrence behaves as $C_\infty(\lambda_c) - C_N(\lambda) = |\lambda - \lambda_c|^{a_{mf}} F_c(N|\lambda - \lambda_c|^\nu)$ with F_c a function of $N|\lambda - \lambda_c|^\nu$ only. When $x \rightarrow \infty$, $F_c(x) \rightarrow \text{constant}$ and $C_\infty(\lambda_c) - C_\infty(\lambda) \sim |\lambda - \lambda_c|^{a_{mf}}$ while for $x \rightarrow 0$, $F_c(x) \rightarrow x^{-a_{mf}/\nu}$ and $C_\infty(\lambda_c) - C_N(\lambda_c) \sim N^{-a_{mf}/\nu}$. In ref. [14] the concurrence in the XY model ($N \rightarrow \infty$, eq. (8)) has been explicitly calculated yielding

$$C(\lambda) = \begin{cases} 1 - \sqrt{1 - \left(\frac{\lambda}{\lambda_c}\right)^2} & , \quad 0 < \lambda < \lambda_c \\ 1 - \sqrt{1 - \left(\frac{\lambda^2}{\lambda_c^2}\right)^2} & , \quad \lambda_c < \lambda < \infty \end{cases} \quad (9)$$

with $\lambda_c = \frac{\sqrt{\epsilon}}{2}$. This implies that, close to λ_c , $1 - C \rightarrow |\lambda - \lambda_c|^{\frac{1}{2}}$. If we assume for the moment that both models, Dicke and infinitely coordinated XY, belong to the same universality class, we can then borrow the result $a_{mf} = \frac{1}{2}$ from the XY concurrence and apply it to the Dicke concurrence. This exponent has also been found for the full Dicke model in ref. [10] using a spin-boson transformation scheme. The exponent $\frac{a_{mf}}{\nu}$ has been numerically obtained (see table I) yielding $\nu = 1.88$. The collapse of all the data on a single curve for each region (coupling below or above the critical coupling strength) is a verification of the existence of scaling behavior of the concurrence in the Dicke model. In fig. 4 (inset) the x-magnetization multiplied by $|\lambda - \lambda_c|^{-1/2}$, is plotted as a function of $N|\lambda - \lambda_c|^{1.43}$. The critical exponent in the argument of the scaling function is 1.43, which is only slightly smaller than that found for the same exponent in the XY model case (1.5). This similarity in scaling exponents reinforces the main message of this paper, that the scaling behaviour of the Dicke model is intimately related to that of the anisotropic infinitely-coordinated XY model.

In summary, we have shown that a radiation-matter (i.e. Dicke) model can be mapped on to an infinitely-coordinated spin/qubit system with an effective magnetic field determined by the detuning between qubit and cavity, and a long-range exchange interaction given by the square of the qubit-cavity coupling strength. An immediate and interesting consequence is that the exchange interaction can be controlled experimentally simply by changing the system's temperature or equivalently the mean number of photons in the cavity. We have also shown that the Dicke model pertains to the same universality class as other infinitely-coordinated systems, such as the anisotropic XY model in a transverse field. We have demonstrated that the finite-size scaling hypothesis works well for the Dicke model, and we therefore postulate that it is also likely to work well for a range of generalized Dicke models corresponding to a diverse range of radiation-matter experimental setups. Temperature-dependent phenomena can also be studied using our size-consistent, effective qubit-qubit system approach.

* * *

We thank A.Olaya-Castro for her valuable comments. This work was supported by the Faculty of Sciences (U. de los Andes), COLCIENCIAS (Colombia) project 1204-05-13614 and DTI-LINK (UK).

REFERENCES

- [1] PAREDES B. ET AL., *Phys. Rev. Lett.*, **90** (2003) 150402.
- [2] PORRAS D. ET AL., *Phys. Rev. Lett.*, **92** (2004) 207901.
- [3] PAREDES B. ET AL., *cond-mat/0306497*, (2003) .
- [4] OSTERLOH A. ET AL., *Nature*, **416** (2002) 608.
- [5] OSBORNE T.J. AND NIELSEN M.A., *Phys. Rev. A.*, **66** (2002) 032110
- [6] VIDAL J. ET AL., *Phys. Rev. A.*, **69** (2004) 022107.
- [7] DICKE R.H., *Phys. Rev.*, **93** (1954) 99.
- [8] WANG Y. K. AND HIOE F. T., *Phys. Rev. A*, **7** (1973) 831.
- [9] EMARY C. AND BRANDES T., *Phys. Rev. Lett.*, **90** (2003) 044101.
- [10] LAMBERT N., EMARY C. AND BRANDES T., *Phys. Rev. Lett.*, **92** (2004) 073602.
- [11] POLATSEK G. AND BECKER K.W., *Phys. Rev. B.*, **55** (1997) 16096.
- [12] WOOTTERS W.K., *Phys.Rev.Lett.*, **80** (1998) 2245.
- [13] BOTET R. AND JULLIEN R., *Phys. Rev.*, **28** (1983) 3955.
- [14] RESLEN J., *Master's Thesis*, edited by UNIVERSIDAD DE LOS ANDES (unpublished) 2004.